



Vision for a Lifetime

Please note:

If you have been using eye drops in the **operative eye** (for Glaucoma, etc) keep taking these drops as prescribed up to the day of surgery BUT **do not use** these drops on the day of surgery. Continue normal routine drops in the non- operative eye.

SHAKE EYE DROPS BEFORE USING

USE DROPS 5 MINTUES APART

EYE DROPS CAN BE USED IN ANY ORDER

VIGAMOX (Pharmacy may substitute Zymar, Quixin, or Iquix)

- Antibiotic

-Start using **after** surgery, use one (1) drop in operative eye three (3) times per day **stop** after two (2) weeks.

NEVANAC (Pharmacy may substitute Xibrom or Acular LS) **or ACUVAIL**

- Non-steroidal anti-inflammatory.

If using Nevanac (or substitution) - Use one (1) drop in operative eye three (3) times per day starting the **day before** surgery. Use the morning of surgery. After surgery continue three (3) times a day **until gone. If diabetic; start using one week before surgery.**

- If using **Acuvail or Xibrom**- Use one (1) drop in operative eye two (**2**) times per day starting the day before surgery. Use the morning of surgery. Continue until gone. **If diabetic; start using one week before surgery.** ****Acuvail** comes in single-use vials.

Omni-Pred (*Sample will be given to you in the clinic. If you did not receive one call 1-800-846-1877 or 651-275-3000*). If samples no longer available will be ePrescribed Omnipred.

- Steroid anti-inflammatory

-Start using **after** surgery, use one (1) drop in operative eye three (3) time per day. Continue until **gone**.

BRING ALL BOTTLES OF EYE DROPS WITH YOU TO THE SURGERY CENTER